



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Am J Surg. 1996 Jan;171(1):192-6.

Related Articles, Link

**ELSEVIER SCIENCE
FULL-TEXT ARTICLE****Adjuvant hormonal treatment with peptide YY or its analog decreases human pancreatic carcinoma growth.****Liu CD, Rongione AJ, Garvey L, Balasubramaniam A, McFadden DW.**

Department of Surgery, UCLA Center for Health Sciences, USA.

BACKGROUND: Recent studies have revealed decreased pancreatic cancer cell growth upon administration of peptide YY (PYY). We examined whether adjuvant treatment with PYY or its synthetic analog, BIM-43004, would decrease human pancreatic adenocarcinoma growth. **MATERIALS AND METHODS:** Human pancreatic ductal adenocarcinomas, MiaPaCa-2 and BxPC-3, were cultured and assessed for growth by MTT assay. Pancreatic cancer cells received 500 pmol of PYY or BIM-43004 for 24 hours prior to 5 fluorouracil (5-FU; 10 micrograms/mL) and leucovorin (40 micrograms/mL) administration. Cell membrane epidermal growth factor (EGF) receptors were analyzed by Western blotting after exposure to peptides and chemotherapy. **RESULTS:** Cancer cell growth was reduced in all groups receiving hormonal pretreatment (23% PYY/5-FU/leucovorin versus control; 27% BIM-43004/5-FU/leucovorin versus control) as compared with groups receiving 5-FU and leucovorin only (16% versus control). The EGF receptor expression was reduced by 30% in cells treated with PYY/5-FU/leucovorin and by 45% in cells treated with BIM/5-FU/leucovorin as compared with control cells without treatment. **CONCLUSION:** Human pancreatic cancer cell growth is further decreased when pretreated with PYY or its synthetic analog prior to chemotherapy.

PMID: 8554139 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Exhibit 2